

Ma cheama si m-am nascut la

Varsta	Locatie	Vaccin	Descriere vaccin obligatoriu	Vaccin optional
0 luni – primele 24h	maternitate	HEP-B	Vaccinare anti-hepatita B	
0 luni – 2-7 zile	maternitate	BCG	Vaccinare anti-tuberculoza	
2 luni	cabinet medical	DTP + VPi + Hib si HEP B (2 administrari)	DTP = vaccinare anti-diftero-tetano-pertussis (impotriva difteriei, tetanosului si tusei convulsive)	Vaccin pneumococic conjugat* (impotriva otitei, meningitei, pneumonii provocate de pneumococ): - daca vaccinarea se incepe la sugar sub 6 luni: 4 administrari la 2, 4, 6, 12 luni - daca vaccinarea se incepe la sugar 6-11 luni: 2 administrari la interval de 1 luna (se recomanda rapel la 12-15 luni) - daca vaccinarea se incepe la 12-23 luni: 2 administrari la interval de 2 luni (fara rapel) -daca vaccinarea se incepe dupa 2 ani: 1 administrare
			VPi = vaccinare anti-polio (impotriva poliomielitei, injectabil)	
			Hib = vaccinare anti-haemophilus influenzae de tip b (provoaca infectii respiratorii dese copiilor mici)	
			HEP B = vaccinare anti-hepatita B.	
4 luni	cabinet medical	DTP + VPi + Hib	DTP = vaccinare anti-diftero-tetano-pertussis (impotriva difteriei, tetanosului si tusei convulsive)	Vaccin anti-rotavirus (impotriva gastroenteritei acute)
			VPi = vaccinare anti-polio (impotriva poliomielitei, injectabil)	
			Hib = vaccinare anti-haemophilus influenzae de tip b (provoaca infectii respiratorii dese copiilor mici)	
6 luni	cabinet medical	DTP + VPi + Hib si HEP B (2 administrari)	DTP = vaccinare anti-diftero-tetano-pertussis (impotriva difteriei, tetanosului si tusei convulsive)	Vaccin anti-rotavirus (impotriva gastroenteritei acute)
			VPi = vaccinare anti-polio (impotriva poliomielitei, injectabil)	
			Hib = vaccinare anti-haemophilus influenzae de tip b (provoaca infectii respiratorii dese copiilor mici)	
			HEP B = vaccinare anti-hepatita B.	
12 luni	cabinet medical	DTP + VPi + Hib	DTP = vaccinare anti-diftero-tetano-pertussis (impotriva difteriei, tetanosului si tusei convulsive)	
			VPi = vaccinare anti-polio (impotriva poliomielitei, injectabil)	
			Hib = vaccinare anti-haemophilus influenzae de tip b (provoaca infectii respiratorii dese copiilor mici)	
13 – 15 luni	cabinet medical	ROR	Vaccinare anti- rubeola, anti-rujeola si anti-oreion	
16 – 18 luni	cabinet medical			Vaccin impotriva hepatitei A (min 3 luni de la ROR) + Vaccin antivaricela
22 – 24 luni	cabinet medical			Vaccin impotriva hepatitei A (rapel la min 6 luni de la prima doza) + Vaccin antimeningococic (f rar)

Ma cheama si m-am nascut la

Varsta	Locatie	Vaccin	Descriere vaccin obligatoriu	Vaccin optional
4 ani	cabinet medical	DTP	Vaccinare anti-diftero-tetano-pertussis (impotriva difteriei, tetanosului si tusei convulsive)	
7 ani (clasa I)	scoala	ROR	Vaccinare anti- rubeola, anti-rujeola si anti-oreion	
9 ani (clasa III)	scoala	VPI	Vaccinare anti-polio (impotriva poliomielitei, injectabil)	
10 – 14 ani	cabinet medical			Vaccin antirubeolic (1 doza, administrat doar fetelor la 10-14 ani sau femeilor neimunizate aflate la varsta fertila)
12 ani	scoala			Vaccinare impotriva papilloma virus (cancer de col uterin)
13 ani (clasa VII)	scoala	BCG (doar pt IDR negativ)	Vaccinare anti-tuberculoza	
14 ani (clasa VIII)	scoala	DT	Vaccinare antidifterica si antitetanica	
24 ani	cabinet medical	DT	Vaccinare antidifterica si antitetanica (ulterior, la fiecare 10 ani)	
Studenti si elevi anul I la facultatile de medicina, stomatologie si scoli sanitare postliceale		HEP-B	Vaccinare anti-hepatita B	